

Supplemental Table 8. Descriptive statistics and ANOVA for craniofacial surgery publications by 3 year periods comparing oromaxillofacial surgery, neurosurgery, and plastic and reconstructive surgery

Time period	Oromaxillofacial surgery		Neurosurgery		Plastic and reconstructive surgery		P-value (ANOVA)
	Mean	SD	Mean	SD	Mean	SD	
2017–2019	58.66	51.68	55.33	3.21	174.00	39.31	<0.001
2014–2016	386.33	37.16	81.33	25.32	327.00	47.84	<0.001
2011–2013	200.00	24.06	32.33	9.45	145.00	33.42	<0.001
2008–2010	128.00	23.38	25.00	4.35	115.00	5.19	<0.001
2005–2007	94.00	7.54	22.00	5.19	79.66	6.02	<0.001
2002–2004	75.33	3.21	15.00	3.60	71.00	7.00	<0.001
1999–2001	79.66	34.12	12.33	2.51	55.00	5.56	0.017
1996–1998	41.66	13.31	12.00	4.35	62.66	15.04	0.006
1993–1995	23.66	1.52	6.00	4.58	39.33	2.51	<0.001
1990–1992	20.66	3.51	4.00	1.00	24.00	6.08	0.002
1987–1989	8.66	5.77	1.00	1.00	11.33	8.62	0.173

SD, standard deviation; ANOVA, analysis of variance.